Garden & Landscape Expo 2020

Orchid Diseases

Brian D. Hudelson
Department of Plant Pathology
University of Wisconsin-Madison/Extension



1





Orchid Diseases Black Rot

- Pathogens
 - Phytophthora cactorum
 - Pythium ultimum
- Favorable environment
 - Excessive moisture
 - Cool temperatures









Orchid Diseases Black Rot

Control

2

4

- Avoid symptomatic plants
- Isolate new plants from established plants
- Decontaminate pots, work surfaces, tools (70% alcohol, disinfectants, 10% bleach)
- Use a proper potting mix
- DO NOT use garden soils

3

Orchid Diseases Black Rot

- Control
 - DO NOT overwater
 - Limit water splash
 - Promote good air circulation
 - Remove diseased plant material and debris

Orchid Diseases Black Rot

- Control
 - Use fungicides to prevent infections
 - · Poultices (?)
 - Copper sulfate + lime
 - Cinnamon
 - Spray treatments
 - Metalaxyl/mefenoxam, fosetyl-Al, etridiazole
 - Mancozeb, captan
 - · Alternate active ingredients (FRAC codes)
 - Apply according to label instructions

5 6

Orchid Diseases Leaf Spots/Blights

Pathogens


- Phyllosticta spp.
- Colletotrichum spp.
- Diplodia spp.
- Uredo spp.

- Septoria spp. - Pseudomonas spp.

Favorable environment

- Excessive moisture
- High humidity

7



8

Orchid Diseases Leaf Spots/Blights

Control

- Avoid symptomatic plants
- Isolate new plants from established plants
- DO NOT get water on leaves
- DO NOT overwater
- Promote good air circulation
- Remove diseased plant material and debris (where possible)

Orchid Diseases Leaf Spots/Blights


Control

- Use fungicides to prevent infections
 - Chlorothalonil, copper, mancozeb, thiophanate-methyl, fludioxonil, captan, ferbam
 - Apply to protect new leaves
 - · 7-14 day application interval
 - Alternate active ingredients (FRAC codes)
 - · Apply during periods of high moisture

9 10

Orchid Diseases Botrytis Blight/Gray Mold

- · Pathogen: Botrytis cinerea
- Favorable environment
 - Excessive moisture
 - High humidity



11 12

Orchid Diseases Botrytis Blight/Gray Mold

- Control
 - Isolate new plants from established plants
 - DO NOT get water on flowers
 - DO NOT overwater
 - Promote good air circulation
 - Remove diseased plant material and debris

Orchid Diseases Botrytis Blight/Gray Mold

Control

14

- Use fungicides to prevent infections
 - Chlorothalonil, copper, fludioxonil, iprodione, mancozeb, maneb, thiophanate-methyl, triflumizole, vinclozolin
 - 7-14 day application interval
 - Alternate active ingredients (FRAC codes)
 - · Apply during periods of high moisture
 - · Apply when "perfect" flowers are desired

13

Orchid Diseases Viruses

- Pathogens
 - Odontoglossum ringspot virus (ORSV)
 - Cymbidium mosaic virus (CymMV)
 - Tobacco mosaic virus (TMV)
 - Cucumber mosaic virus (CMV)
 - Impatiens necrotic spot virus (INSV)
 - Tomato spotted wilt virus (TSWV)
 - Others

Orchid Diseases

Viruses

· Favorable environment: None

Transmission

- Touch: TMV

- Mechanical: ORSV, CymMV

- Aphid: CMV

- Thrips: INSV, TSWV

15 16



Orchid Diseases Viruses

• Control

- Avoid symptomatic plants
- Isolate new plants from established plants
- Request testing of plants prior to purchase
 - Enzyme-Linked Immunosorbent Assay (ELISA)
 - · Dip-stick tests
 - Polymerase Chain Reaction (PCR)
- Discard infected plants

17 18

Orchid Diseases

Viruses

Control

- Dinfest contaminated materials
 - 1% sodium dodecyl sulfate (sodium lauryl sulfate)
 - + 1% Alconox® (2½ Tbsp + 2¾ Tbsp/gal)
 - Trisodium phosphate (14 dry oz/gal)
 - 20% low fat dry milk (Carnation ®) + 0.1% polysorbate 20 (91% cups + 3/4 tsp/gal)
 - · Alcohol dip followed by flaming

Orchid Diseases Where to Go for Help

Plant Disease Diagnostics Clinic
Department of Plant Pathology
University of Wisconsin-Madison
1630 Linden Drive
Madison, WI 53706-1598
(608) 262-2863
pddc@wisc.edu
https://pddc.wisc.edu
Follow on Facebook and Twitter @UWPDDC