Wisconsin Hosta Society

Hosta Diseases

Brian D. Hudelson
Department of Plant Pathology
University of Wisconsin-Madison/Extension







Hosta Diseases

Hosta Virus X

• Pathogen: Hosta virus X (HVX)

· Host: Hosta

- 'Gold Edger' - 'Striptease'

- 'Gold Standard' - 'Sum and Substance'


- 'Golden Tiara'

· Favorable environment: None

Hosta Diseases

Hosta Virus X

- · Transmission: Mechanical
 - Shovels
 - Knives
 - String edgers
 - Lawnmowers



Hosta Diseases

Hosta Virus X

- Control
 - Buy hostas from a reputable source
 - · Inspect hostas for symptoms
 - · Test hostas prior to purchase
 - Keep new and established hostas separated
 - Remove diseased plants and plant debris
 - · Burn (where allowed)
 - Deep bury
 - Hot compost

Hosta Diseases Hosta Virus X

- Control
 - Disinfest contaminated materials
 - 1% Sodium dodecyl sulfate (sodium lauryl sulfate) + 1% Alconox® (2½ Tbsp + 2¾ Tbsp/gal)
 - Trisodium phosphate (14 dry oz/gal)
 - 20% low fat dry milk (Carnation®) + 0.1% polysorbate 20 (91% cups + 34 tsp/gal)
 - · Alcohol dip followed by flaming
 - DO NOT use chemical controls on plants

Hosta Diseases

Tobacco Rattle


- Pathogen: Tobacco rattle virus (TRV)
- Hosts
 - Herbaceous ornamentals
 - Astilbe, bleeding heart, columbine, coral bells, daffodils, epimedium, gladiolus, hosta, hyacinth, marigold, peony, tulip, vinca
 - Vegetables
 - · Beans, beet, pepper, potato, spinach
- · Favorable environment: None

Hosta Diseases

Tobacco Rattle

- Transmission
 - Stubby-root nematodes
 - · Trichodorus spp.
 - · Paratrichodorus spp.
 - Mechanical
 - Grafting
 - Seed





Hosta Diseases Tobacco Rattle

- Control
 - DO NOT buy symptomatic plants
 - Grow non-susceptible plants
 - Annual phlox, carnation, devil's trumpet (downy thorn-apple), sweet William, zinnia, zombie cucumber
 - Remove and destroy infected plants
 - · Burn (where allowed)
 - Deep bury
 - · Hot compost

Hosta Diseases

Tobacco Rattle

- Control
 - Disinfest contaminated materials
 - 1% Sodium dodecyl sulfate (sodium lauryl sulfate) + 1% Alconox® (2½ Tbsp + 2¾ Tbsp/gal)
 - Trisodium phosphate (14 dry oz/gal)
 - 20% low fat dry milk (Carnation®) + 0.1% polysorbate 20 (9% cups + 3/4 tsp/gal)
 - · Alcohol dip followed by flaming

Hosta Diseases


Tobacco Rattle

- Control
 - DO NOT use chemical controls on plants
 - DO NOT attempt to control stubby-root nematodes

Hosta Diseases

Foliar Nematode

- Pathogen: Aphelenchoides spp.
- Hosts
 - Many types of herbaceous plants
 - Landscape plants
 - · Hosta, begonia, coral bells, miterwort
 - Houseplants
 - · African violets, ferns, chrysanthemum
- Favorable environment: Wet weather





Hosta Diseases Foliar Nematode

- Control
 - Inspect plants prior to purchase
 - Avoid overhead watering
 - Remove symptomatic plants and plant debris
 - Burn (where allowed)
 - · Deep bury (landfill)
 - Hot compost

Hosta Diseases Foliar Nematode

- Control
 - Disinfest contaminated materials
 (70% alcohol, disinfectants, 10% bleach)
 - Hot water treatments (10 minutes at 125°F) $\,$
 - DO NOT use nematicides

Hosta DiseasesSouthern Blight

• Pathogen: Sclerotium rolfsii

Hosts
 Hosta

- Many bedding plants

- Many herbaceous annuals and perennials

- Some woody ornamentals

· Favorable environment: Warm, wet weather



Hosta Diseases Southern Blight

- Control
 - DO NOT buy infected/infested plants
 - Avoid cocoa mulch (?)
 - Remove infected plants, mulch and soil
 - Double bag and landfill
 - Disinfest contaminated materials
 (70% alcohol, disinfectants, 10% bleach)

Hosta Diseases Southern Blight

- Control
 - Amend soil with organic matter (?)
 - Use fungicides for control
 - Contract with a professional pesticide applicator
 - Azoxystrobin, flutolanil, flutolanil + thiophanatemethyl, PCNB, tebuconazole, triadimefon
 - · Alternate active ingredients (FRAC codes)
 - Apply at 14 28 day intervals
 - Pray for a really, really, REALLY cold winter

Hosta Diseases Root/Crown Rots

- Pathogens
 - Pythium spp. Phytophthora spp.
 - Rhizoctonia solani Fusarium spp.
 - Cylindrocarpon spp. Thielaviopsis spp.
- Hosts
 - Hosta
 - Any herbaceous plant
- · Favorable environment: Cool, wet soils



Hosta Diseases Root/Crown Rots

- Control
 - Moderate soil moisture
 - · Grow plants in well-drained sites
 - · Use a potting mix with adequate drainage
 - · Improve drainage in poorly drained soils
 - Add organic matter to improve drainage
 - Use raised beds
 - DO NOT overwater
 - DO NOT overmulch

Hosta Diseases Root/Crown Rots

- Control
 - Pretest soils/mulches/composts
 - Use a soil-less potting mix or a pasteurized potting mix for containerized plants
 - DO NOT move contaminated soil or plants
 - Disinfest contaminated materials
 (70% alcohol, disinfectants, 10% bleach)

Hosta Diseases Root/Crown Rots

- Control
 - Use fungicides to prevent infections
 - · Contract with a professional pesticide applicator
 - Etridiazole, metalaxyl, mefenoxam, fosetyl-Al, PCNB, thiophanate-methyl, fludioxonil
 - · Alternate active ingredients (FRAC codes)
 - · Use granular formulations if possible
 - · Use during periods of wet weather

Hosta Diseases Root/Crown Rots

- Control
 - Use biopesticides to prevent infections
 - · Trichoderma, Gliocladium
 - Use for potted plants
 - Exercise caution when combining with chemical fungicides

Hosta Diseases Sun Scald/Sunburn

- Causes
 - -Excessive light
 - -Excessive heat
 - -Insufficient rain
- Hosts
 - -Hosta
 - -Other shade-loving plants



Hosta Diseases

Sun Scald/Sunburn

- Management
 - Site plants appropriately
 - Provide adequate shade
 - Water adequately
 - Mulch appropriately

Hosta Diseases Where to Go for Help

Plant Disease Diagnostics Clinic
Department of Plant Pathology
University of Wisconsin-Madison
1630 Linden Drive
Madison, WI 53706-1598
(608) 262-2863
pddc@wisc.edu
https://pddc.wisc.edu
Follow on Facebook and Twitter @UWPDDC

Subscribe to the PDDC Listserv: UWPDDCLearn